


A



B




Figure S1. Protein secretion from IFN- β treated, St infected macrophages and CXCL2 induction in the Ileal Epithelial MODE-K cell line.

(A) WT C57BL6/J macrophages were pretreated for 60 minutes with media alone or 100 u/ml murine IFN- β and infected with St strain SL1344 for 24 hours at an moi 4. Cell supernatants were harvested and chemokine and cytokine levels analyzed by ELISA. (B) 4×10^5 MODEK murine epithelial cells were plated per well. Cells were pre-treated for 4 hours with media alone or media supplemented with 100 U/ml IFN β . Cells were subsequently infected with SL1344 at an moi of 4 for 6 hours and mRNA levels of CXCL2 assayed by qRT-PCR.




Figure S2. Protein Secretion from IFN- $\beta^{+/+}$ and IFN- $\beta^{-/-}$ St infected macrophages. WT or IFN- $\beta^{-/-}$ macrophages were infected with St strain SL1344 for 24 hours at an moi 4. Cell supernatants were harvested and chemokine and cytokine levels analyzed by ELISA




Figure S3. IFN- β does not suppress TLR4 expression, Intracellular Trafficking, or Signal Transduction

(A) WT C57BL6/J peritoneal macrophages were treated for 6 hours with IFN- β and steady state expression levels of the indicated transcripts was determined by qRT PCR. (B) WT C57BL6/J peritoneal macrophages were treated as in (A) and steady state surface expression of TLR4 was quantified by Flow cytometry. (C) Media or IFN- β pre-treated macrophages were stimulated with 100 ng/ml purified *E. coli* LPS. At the indicated times, surface TLR4 was assayed by flow cytometry. The remaining amount of surface TLR4 is expressed as a percent of MFI from non LPS treated cells. (D) Media or IFN- β pre-treated macrophages were stimulated with 100 ng/ml *E. coli* LPS for 30 minutes and whole cell lysates were probed with antibodies against the indicated proteins by western blot. (E) WT C57BL6/J and TLR4^{-/-} macrophages were treated for 4 hours with media alone or with 100 U/ml IFN- β and subsequently infected for 2 hours with SL1344 at an moi of 4. Whole cell lysates were probed by western blot with the indicated antibodies