



**Supplemental Figure 1: TLR-dependent production of IFN- $\gamma$  by T cells** (A) Intracellular IFN- $\gamma$  staining was performed on MACS-enriched CD8<sup>+</sup> OT-I T cells directly post selection (*ex vivo*), after activation with MEC.B7.SigOVA cells for 20h or at indicated time points of rest in the absence of antigen. T cells were left untreated, or were incubated for 6h with 1 $\mu$ g/ml brefeldin A (BFA), in the presence or absence of 1 $\mu$ g/ml OVA<sub>257-264</sub> peptide or the irrelevant SV40<sub>560-568</sub> peptide. Numbers in the upper right corner indicate the percentage of IFN- $\gamma$ <sup>+</sup> CD8<sup>+</sup> T cells. (B) MACS-selected T cells (rested for 3 days) were pre-incubated for 30 min with 10 $\mu$ g/ml IL-12 neutralizing antibody or left untreated (-), and then stimulated for 6h with Pam3 and R848 in the presence or absence of the IL-12 neutralizing antibody. IFN- $\gamma$  production was measured by flow cytometry. (C-D) FACS-sorted naive CD44<sup>low</sup> CD62L<sup>hi</sup> CD8<sup>+</sup> OT-I T cells were activated with MEC.B7.SigOVA cells for 20h and removed from antigenic stimuli for indicated time points. (C) Dot plots represent IFN- $\gamma$  and TNF- $\alpha$  (upper panel), and IFN- $\gamma$  and IL-2 production (lower panel) of T cells (rested for 3 days) that were stimulated for 6h with 5 $\mu$ g/ml Pam3 or 10 $\mu$ g/ml R848. (D) Intracellular IFN- $\gamma$  staining of T cells that were activated with Pam3 and R848 upon 6 days, or 15 days of rest.



**Supplemental Figure 2: Pam3 stimulation of IFN- $\gamma$ R<sup>-/-</sup> macrophages.** BM derived-macrophages from Balb-c WT and IFN- $\gamma$ R<sup>-/-</sup> mice were cultured for 5h with 5 $\mu$ g/ml Pam3 or left untreated. (A) TNF- $\alpha$  and (B) IL-10 mRNA expression was analyzed by RT-PCR. Data are pooled from 2 independently performed experiments.



**Supplemental Figure 3: TCR-dependent IFN- $\gamma$  production is controlled by both mTORC1 and mTORC2 pathways, and is supported by glycolysis** (A) Dot plots represent Akt, mTOR and S6 phosphorylation 30 minutes after T cell reactivation with 5 $\mu$ g/ml Pam3 or 100nM OVA<sub>257-264</sub> peptide. When indicated, cells were pretreated for 30 minutes with ZTS inhibitor. (B) Intracellular IFN- $\gamma$  staining of CD8<sup>+</sup> T cells that were activated for 6h with 5 $\mu$ g/ml Pam3, 0.1nM, 1nM or 100nM OVA<sub>257-264</sub> peptide or were left unstimulated. (C) Resting T cells were incubated for 30 min with indicated concentrations of rapamycin or left untreated prior to reactivation with 1nM OVA<sub>257-264</sub> peptide. Graph indicates percentage of IFN- $\gamma$  producing cells relative to cells that were reactivated in the absence of rapamycin. Data are pooled from 4 independent experiments  $\pm$  SD [one-way ANOVA with Dunnett's multiple comparison to untreated control; \*\*p<0.005; \*\*\*\*p<0.0001]. (D) Akt, mTOR and S6 phosphorylation of T cells reactivated for 30 minutes with low amount (1nM) of OVA<sub>257-264</sub> peptide. (E) T cells were activated for 6h with 1nM OVA<sub>257-264</sub> peptide in FCS-free, glucose-free RPMI that was supplemented with 25mM glucose, or with 25mM glucose plus 10mM 2-DG, or left unsupplemented. [one-way ANOVA with Dunnett's multiple comparison to glucose condition; \*\*p<0.005]. (F) Glucose-dependency, (G) glutamine-dependency and mitochondrial respiration-dependency assays were performed when T cells were rested for 15 day and then stimulated for 6h with 5 $\mu$ g/ml Pam3 or 100nM OVA<sub>257-264</sub> peptide. [one-way ANOVA with Dunnett's multiple comparison to negative condition; \*p<0.05; \*\*p<0.005].

Supplemental Table I: Primers used for RT-PCR

<b>Gene (mouse)</b>	<b>Forward sequences</b>	<b>Reverse sequences</b>
L32	5'-GAAACTGGCGGAAACCCA-3'	5'-GGATCTGGCCCTTGAACCTT-3'
IFN- $\gamma$	5'-GGATGCATTCATGAGTATTGC-3'	5'-CCTTTTCCGCTTCCTGAGG-3'
TNF- $\alpha$	5'-ACAGAAAGCATGATCCGCG-3'	5'-GCCCCCATCTTTTGGG-3'
IL-10	5'-GGTTGCCAAGCCTTATCGGA-3'	5'-ACCTGCTCCACTGCCTTGCT-3'
TLR2	5'-AAGAGGAAGCCCAAGAAAGC-3'	5'-CGATGGAATCGATGATGTTG-3'
TLR3	5'-GGTGGTCCCGTTAATTCCT-3'	5'-CCCGAAAACATCCTTCTCAA-3'
TLR4	5'-TCAGAACTTCAGTGGCTGGATTT-3'	5'-AACTCTGGATAGGGTTTCTGTC-3'
TLR5	5'-AAGTTCCGGGGAATCTGTTT-3'	5'-GCATAGCCTGAGCCTGTTTC-3'
TLR7	5'-GGCTGAACCATCTGGAAGAA-3'	5'-TAAGCTGGATGGCAGATCCT-3'
TLR9	5'-AGCTGAACATGAACGGCATCT-3'	5'-CCAGCCATCTGAGCGTGTACT-3'